



MATRIXYL®3000

Patent N° WO 2005/048968

matrixyl
inside



Pal-GHK and Pal-GQPR

Function:

Anti-wrinkle and lifting effect.

Definition:

Matrixyl®3000 contains two matrikines®, Pal-GHK and Pal-GQPR which act in synergy to restore and maintain skin's youthful appearance.

Properties:

Matrixyl®3000 contains matrikines which are messengers of cutaneous restructuring and repair. They activate the neosynthesis of extracellular matrix macromolecules providing Matrixyl®3000 with a visible anti-wrinkle efficacy.

Characteristics:

As messenger molecules, matrikines® are capable of regulating cell activities. They interact with specific receptors to activate certain genes involved in the process of extracellular matrix renewal and cell proliferation. With age these mechanisms become progressively weaker.

INCI Name:

(Check CTFA on-line dictionary for latest INCI name)
Glycerin – Aqua (Water) – Butylene Glycol – Carbomer – Polysorbate 20 – Palmitoyl Oligopeptide – Palmitoyl Tetrapeptide-7*
* former INCI name: Palmitoyl Tetrapeptide-3

Applications:

Anti-aging , face lifting and wrinkle treatment products. Suitable for gels, serums lotion, creams.

Formulation:

Water soluble.

Recommended use level:

3% to 8%

45%
less wrinkles
in 2 months



Lifetime youth



www.sederma.fr
E-mail: sederma@sederma.fr
Copyright© 2000-2010
Sederma. All rights reserved.

Member of Croda International Plc


In vitro tests

● SYNTHESIS OF MATRIX MACROMOLECULES

Study to determine ability of fibroblasts incubated for 72 hours with Matrixyl®3000 (1, 3, and 5%) to stimulate synthesis of extracellular matrix components.

● STIMULATION OF GENE EXPRESSION

Study to evaluate regulation of dermal and epidermal genes by matrikines present in Matrixyl®3000, using DNA-Array techniques on a reconstructed epidermis and a fibroblast culture.



The profile of gene activation by the matrikines® in Matrixyl®3000 complement the skin's natural mechanisms for cutaneous restructuring.

In vivo tests

● ANTI-WRINKLE EFFICACY

Clinical study in which 23 volunteers aged 39 to 74 applied a cream containing 3% Matrixyl®3000 to one-half of their face against a placebo on the other half, twice a day for 56 days. Anti-wrinkling and lifting efficacy was assessed by profilometry and photography, as compared to T0.



VARIATION OF PARAMETERS COMPARED TO T0 (%)

	Matrixyl®3000	Placebo
Surface occupied by deep wrinkles (>200µm)	- 44.9**	4.3ns
Main wrinkle density	- 37.0**	- 9.6ns
Main wrinkle average depth	- 15.1**	- 3.2ns
Main wrinkle average volume	- 18.5**	- 8.7*
Roughness	- 14.4**	1.4ns
Complexity (Lifting effect)	- 16.6**	4.2ns

ns : non significant *significant / T0 (p<0.05) **significant / T0 (p<0.01)

● EFFECT ON SKIN TONE AND ELASTICITY

Assesment of elasticity and tone by cutometry on the same panel as above.



Matrixyl®3000 displays major and significant improvements in elasticity and tone.